

ADDITIONS TO "AN ANNOTATED LIST OF SOUTH CAROLINA SALTICIDAE (ARANEAE)"

S. H. Roach and G. B. Edwards

Cotton Production Research Unit, USDA/ARS, P. O. Box 2131, Florence, SC 29503; and
Florida State Collection of Arthropods, P. O. Box 1269, Gainesville, FL 32602, respectively.

Several species of salticids have been added to the list of species previously reported as occurring or possibly occurring in South Carolina since publication of our original list (Roach and Edwards 1984). The following is a list of species and collection data for addition to the previously published list.

Admestina tibialis (C. L. Koch). One adult female collected from a retreat on ground moss in Pickens County, South Carolina, June 1985, by J. Brushwein.

Peckhamia americana (Peckham & Peckham). One adult female collected 16 May 1986, in mixed hardwood, Darlington County, South Carolina, by SHR.

Phidippus apacheanus Chamberlin & Gertsch. One adult male from a fall armyworm trap in Florence County, South Carolina, on 23 September 1986. The trap was located on a ditch bank between fields of tobacco and corn. Another male was collected in Long Bay, Myrtle Beach, Horry Co., South Carolina, on 15 October 1940, by E. Mayr (GBE).

Phidippus cardinalis (Hentz). One adult male collected from shrubs, Florence County, South Carolina, on 25 September 1986, by SHR.

Sassacus papenhoei Peckham & Peckham. One female d-vac collected in cotton, Society Hill, Darlington Co., South Carolina, 25 June 1987, by SHR. This species is extremely rare in the eastern U.S. (GBE).

Sitticus magnus Chamberlin & Ivie. One male and one female collected in mixed hardwood on low vegetation, Florence, Florence Co. (at Darlington Co. line), South Carolina, May 1986, by SHR.

Thiodina peurpera (Hentz). One adult male collected from shrubs, Florence County, South Carolina, 25 September 1986, by SHR.

Brimley (1938) listed the following species collected by N. Banks in North Carolina which may also occur in the mountainous areas of South Carolina:

Hasarius adansonii (Audouin). From Mt. Graybeard.

Metaphidippus insignis Banks. From Roan Mountain.

Brimley also noted in the same list *Sassacus papenhoei*, from Blowing Rock. We report above this species from South Carolina for the first time.

REFERENCES

- Brimley, C. S. 1938. Near Insects, pp. 467-503, *in*, Insects of North Carolina. N. Car. Dept. Agric., Div. Entomol., Raleigh, NC. 560 pp.
- Roach, S. H. and G. B. Edwards. 1984. An Annotated List of South Carolina Salticidae (Araneae). *Peckhamia* 2: 49-56.